
# CHROMagar™ Mastitis (GN)



# CHROMagar™ Mastitis (GN)

### **MEDIUM PURPOSE**

For isolation and differentiation of the main gram negative pathogens involved in Mastitis infections.

# **COMPOSITION**

The product is composed of a single powder base.

Product =	Base
Total g/L	33.2 g/L
Composition g/L	Agar 15.0 Peptone and yeast extract 17.0 Chromogenic mix 1.2
Aspect	Powder Form
STORAGE	15-30 °C
FINAL MEDIA pH	7.0 +/- 0.2

## PREPARATION (Calculation for 1L)

# Step 1 Preparation of the mix

- Disperse slowly 33.2 g of powder base in 1 L of purified water.
- Stir until agar is well thickened.
- Heat and bring to boil (100 °C) while swirling or stirring regularly.

Advice 1: For the 100°C heating step, mixture may also be brought to a boil in a microwave oven: after initial boiling, remove from oven, stir gently, then return to oven for short repeated bursts of heating until complete fusion of the agar grains has taken place (large bubbles replacing foam).

DO NOT HEAT TO MORE THAN 100 °C. DO NOT AUTOCLAVE AT 121 °C.

# Step 2 Pouring

- Cool in a water bath to 45-50 °C, swirling or stirring gently.
- Pour into sterile Petri dishes.
- Let it solidify and dry.

### Storage

- Store in the dark before use.
- Prepared media plates can be kept for one day at room temperature.
- $\bullet$  Plates can be stored for up to 1 month under refrigeration (2/8 °C) if properly prepared and protected from light and dehydration.

### **INOCULATION**

Related samples can be processed by direct streaking on the plate, as well as prior appropriate enrichment step.

- $\bullet \ \ \text{If the agar plate has been refrigerated, allow to warm to room temperature before inoculation}.$
- Streak sample onto plate.
- Incubate in aerobic conditions at 37 °C for 18-24 hours.

### **Typical Samples**

e.g. milk

Possible enrichment step Direct streaking or spreading technique


**ENGLISH** 

# CHROMagar™ Mastitis (GN)

### **INTERPRETATION**

Microorganism	Typical colony appearance
E.coli	ightarrow dark pink to reddish
Klebsiella, Enterobacter, Citrobacter	→ metallic blue (+/- reddish halo)
Proteus, Morganella, Providencia	→ brown halo
Pseudomonas	→ translucent ( +/- natural pigmentation cream to green)
Candida albicans	→ cream, pinpoint colonies
Gram positive bacteria	→ inhibited

## <u>Typical</u> colony appearance







The pictures shown are not contractual.

### PERFORMANCE & LIMITATIONS

• Final identification may require additional tests such as indole test (on paper), ONPG, Oxidase.

### **QUALITY CONTROL**

√Σ∕ Pack Size

5000 ml

Please perform Quality Control according to the use of the medium and the local QC regulations and norms.

Good preparation of the medium can be tested, isolating the ATCC strains below:

Microorganism	Typical colony appearance
K.pneumoniae ATCC® 13883	→ metallic blue
E.coli ATCC® 25922	→ reddish
P.aeruginosa ATCC® 10145	→ translucent
S.aureus ATCC® 25923	→ inhibited
E.faecalis ATCC® 29212	→ inhibited

**Ordering References** 

MS252/N

### **WARNINGS**

- Do not use plates if they show any evidence of contamination or any sign of deterioration.
- Do not use the product beyond its expiry date or if product shows any evidence of contamination or any sign of deterioration.
- For Laboratory use. This laboratory product should be used only by trained personnel in compliance with good laboratory practices.
- Any change or modification in the procedure may affect the results.
- Any change or modification of the required storage temperature may affect the performance of the product.
- Unappropriate storage may affect the shelf life of the product.
- Recap the bottles tightly after each preparation and keep them in a low humidity environment, protected from moisture and light.
- For a good microbial detection: collection and transport of specimen should be well handled and adapted to the particular specimen according to good laboratory practices.

### **DISPOSAL OF WASTE**

After use, all plates and any other contaminated materials must be sterilized or disposed of by propriate internal procedures and in accordance with local legislations. Plates can be destroyed by autoclaving at 121°C for at least 20 minutes.

#### **REFERENCES**

Please refer to our website page «Publications» for scientific publications about this particular product.

Web link: http://www.chromagar.com/publication.php

### IFU/LABEL INDEX

Σ

Quantity of powder sufficient for X liters of media



Expiry date



Required storage temperature



Base (B)

MS252/N

Weight: 166 gr

Store away from humidity

Need some Technical Documents?

Available for download on www.CHROMagar.com

- Certificate of Analysis (CoA) --> One per Lot
- Material Safety Data Sheet (MSDS)

CHROMagar  $^{\text{TM}}$  and Rambach  $^{\text{TM}}$  are trademarks created by Dr A. Rambach ATCC $^{\circ}$  is a registered trademark of the American Type Culture Collection NT-EXT-097 V1.0 / 04-Apr-17



250 Tests

of 20ml

CHROMagar 4 place du 18 juin 1940 75006 Paris - France Email: CHROMagar@CHROMagar.com Tel +33 (0)1.45.48.05.05. Website: www.CHROMagar.com